

Marphysa sp. A station	width	branchiae on setigers#	average # branchial filaments	capillary setae	composite setae	occipital tentacles	eyes	prostomium	acicula color & number	subacicular hooks	pectinate setae	comments
LBGS-NPDES Bio- specimen 1 	1 mm	8/10 thru 16/18	8		falcigers present in posterior				black			possibly young M. disjuncta
LBGS-NPDES Bio- specimen 2 	1.75 mm	11 thru 20	10-11		falcigers present in posterior				black			possibly young M. disjuncta
LBGS-NPDES Bio- specimen 3 	3 mm	12 thru 26/29	16-17		falcigers present in posterior				black			possibly young M. disjuncta
LBGS-NPDES Bio- specimen 4 	4.5 mm	13 thru 29	19-20		falcigers present in posterior				black			possibly young M. disjuncta
Goleta & Encina		7 thru 13	5-6		falcigers & spinigers in post-branchial region				black			
Catalina Canyon, 379 m (Hartman 1963, 1968)	1.1 mm	10 thru 21	up to 8	present	falcigers & spinigers in post-branchial region	longest extends to 2, black setiger 2, smooth		rounded	pale at tips, dark basally, distally blunt	bifid, ill. shows rounded tips	present	Hartman (1968) identified this as M. belli oculata Treadwell 1921; her description is of a small specimen of sp A, some characters are size-related