

TABLE 1

Table of systematic criteria in *Listriella* from southern California

		color	pigment head cap	antennae long	eyes present	pleon 3 notched	uropod 3 outer ramus=inner ramus	per. 7, art 7 long	per. 7, art. 6 paddle shaped	uropod 1 with ped. spine	ant. 1 article 2 has pigment	male-female uropod 3 is the same
<u>melanica</u>	MALE	K	O	O	X	X	X	X	O	X	X	X
	FEMALE	K	O	O	X	X	X	X	O	X	X	
	JUVENILE	K	O	O	X	X	X	X	O	X	X	
<u>Diffusa</u>	MALE	D	O	O	X	X	X	X	OX	X	O	O
	FEMALE	D	O	O	X	X	X	X	OX	X	O	
	JUVENILE	D	O	O	X	X	X	X	OX	X	O	
<u>goleta</u>	MALE	DS	H	X	X	X	X	X	O	X	X	X
	FEMALE	DS	H	XO	X	X	X	X	O	X	X	
	JUVENILE	DS	H	O	X	X	X	X	O	X	X	
<u>eriopisa</u>	MALE	S	X	O	X	O	O	O	X	XO	O	X
	FEMALE	S	X	O	X	O	O	O	X	XO	O	
	JUVENILE	S	X	O	X	X	XO	OX	XO	XO	O	
<u>albina</u>	MALE	O	O	OX	O	X	XO	X	OX	X	O	X
	FEMALE	O	O	OX	O	X	XO	X	OX	X	O	
	JUVENILE	O	O	OX	O	X	XO	X	OX	X	O	
sp. A	MALE	KS	O	O	X	O	X	X	O	O	X	X
	FEMALE	KS	O	O	X	O	X	X	O	O	O	X
	JUVENILE											

K = dark  
 S = striped  
 DS = diffused-striped  
 X = positive  
 O = negative  
 OX and XO = intermediate  
 H = hourglass-shaped mark

(modified from Barnard, 1959)