

Campylaspis rubromaculata Lie 1969
Nannastacidae

SCAMIT Vol. 4, No. 12

SCAMIT Code: LACO 67

Date Examined: Feb. 10, 1986
Voucher by: Douglas Diener

Synonymy: Campylaspis (papillata) lomikina in Lie 1968
Campylaspis nodulosa Lie, 1969
Campylaspis rubromaculata Lie, 1971

Literature: Lie, U. 1968, 1969, 1971

Diagnostic Characters

1. Generally heavily pigmented (orange to red).
2. Carapace with two complete and one partial lateral ridge (see figure).
3. Inner margin of uropodial peduncle with 4 to 6 small spines.

Related Species and Character Differences:

Campylaspis rubromaculata can be confused with C. hartae; compare the carapace sculpturing pattern carefully.

Distribution: Puget Sound to San Diego; sand, silt bottoms, kelp hold fast; 6-640 m.






Figure 1. Carapace (from Lie, 1969)

Figure 2. Telson (from Lie, 1969)

Figure 3. Carapace (from MBC, unpublished)

Campylaspis
rubromaculata Lie, 1971
Nannastacidae



SUPPLEMENTAL FIGURES

SCAMIT Vol. 6, No. 12


SCAMIT Code: LACO 67 (in part)
MEC 2
Also specimens DL-SONGS

Date Examined: January 11, 1988
Voucher By: Doug Diener, MEC

Voucher Sheet: Vol. 4, No. 12



Female



Male