

A KEY TO CORYMORPHINE POLYPS

1. Both whorls filiform to submoniliform, tips bulbous: papillae at base of hydrocaulus
.....*Corymorpha* 2
1. Aboral whorl moniliform, capitate; oral whorl capitate, sub- to moniliform; papillae at top of hydrocaulus, below line of demarcation.....*Euphypha* 3
2. Gonangia are cryptomedusae (elongate, fusiform bodies)*Corymorpha palma*
2. Gonangia are quadrate eumedusoids with one tentacle longer...*Corymorpha bigelowi*
2. Very short hydrocaulus, equal to or shorter than hydranth....*Corymorpha* sp. A
3. Oral whorl tentacles profuse, long, sub- to moniliform, capitate; quadrate eumedusoids with 4 equal tentacles*Euphypha* sp. B
3. Oral tentacles 4-9, short and capitate; hydrocaulus tapering : hypostome short and blunt; about 1mm.....*Euphypha* sp. A
3. Oral tentacles 3-7, short and capitate; hydrocaulus thin with uniform diameter; hypostome ovoid.....*Euphypha ruthae*

CORYMORPHINES

- Brinckmann-Voss, A. 1970. Anthomedusae/ Athecatae (Hydrozoa, Cnidaria) of the Mediterranean. Part I. Capitata. Flora Fauna Golfo Napoli, 39: 1-36.
- Campbell, R.D. 1968. Holdfast movement in the hydroid *Corymorpha palma*: a mechanism of elongation. Biol. Bull. (Woods Hole), 134:26-34.
- Iwasa, M. 1934. Revision of *Stylactis* and its allied genera, with description of *Stylactella (Stylactis) yerii* n.sp. J. Fac. Hokkaido Univ. Ser. 6, Zool.; Vol. 2 no. 4, pp. 241-277.
- Kramp, P.L. 1949. Origin of the Family Corymorphidae. Vidensk. Medd. Dan. Naturhist. Foren., 107: 13-32.
- Miles, S.S. 1937. A new genus of hydroid and its method of asexual reproduction. Biol. Bull. (Woods Hole), 72: 327-333.
- Murbach, L. 1899. Hydroids from Woods Hole, Mass. *Hypolytus peregrinus*, , a new unattached marine hydroid: *Corynitis Agassizii* and its medusa. Q.J. Microsc. Sci., 42: 341-360.
- Rees, W.J. 1937. On the systematic position of the hydroid genera *Hypolytus* Murbach, 1899 and *Dahlgrenella* Miles, 1937. Ann. Mag. Nat. Hist., 13: 102-106.
- Rees, W.J. 1957. Evolutionary trends in the classification of capitate hydroids and medusae. Bull. Br. Mus. N. Hist., Zoology, Vol. 4, no. 9, pp. 455-534.
- Naumov, D.V., 1969. Hydroids and Hydromedusae of the USSR. Israel Program for Scientific Translations, Jerusalem. 660 pp.
- Norenburg, J.L., and M.P. Morse. 1983. Systematic implications of *Euphypha ruthae* n.sp. (Atheata: Corymorphidae), a psammophilic solitary hydroid with unusual morphogenesis. Trans. Amer. Microsc. Soc., Vol. 102, no. 1, pp. 1-17.
- Petersen, K.W. 1990. Evolution and taxonomy in capitate hydroids and medusae (Cnidaria: Hydrozoa). Zool. J. Linn. Soc. Vol. 100. pp 101-231.
- Sassaman, Clay and J.T. Rees. 1978. The life cycle of *Corymorpha* (=*Euphypha*) *bigelowi* (Maas, 1905) and its significance in the systematics of Corymorphid hydromedusae. Biological Bulletin, Vol 154: 485-496.
- Torrey, H.B. 1907. Biological studies on *Corymorpha*. II. The development of *C. palma* from the egg. Univ. Cal. Publ. Zool., 3 (12): 253-298.