

Voucher #


November 14, 1983

Literature Citation: Hinds, 1845: pg 67; plate 21, figure 2

Synonymy: Tellina santarosae Dall, 1900

Primary Diagnostic Characters: Medium-sized (to 60mm), elongate; flattened; adults thick; longer, rounded anteriorly; pointed, slightly truncate posteriorly; sculpture of heavy, closely spaced concentric ribs; white, sometimes with a slight yellow or pink hue internally (Coan, 1971).

Variability: Southern specimens of this species differ, on an average, from northern material, generally flatter shape, thinner shell, and minor sculptural differences (Burch, 1945a).



from Fitch 1953





Figure 2
Tellina bodegensis, internal view of valves
from Coan 1971



Pertinent Literature:

I. Oldroyd, 1925: pg. 168; plate 44, figure 5.

Keen, 1966b: pg. 267

Coan, 1971: pp. 10-11, plate 1, figures 4,5

Depth Range: 0-96m

Distribution:

Burch 1945, Queen Charlotte Island, B.C. to Gulf of California. Japan?

Abbott 1974, Graham Island, British Columbia, to the Gulf of California.

Fitch 1953, Queen Charlotte Island, B.C. to Cape San Lucas, Baja California

Ecology:

Bottom type: sandy sediment on exposed beaches, moderately common