

SIPUNCULA

by

BRUCE E. THOMPSON

SCCWRP

646 W. P.C.H.

Long Beach, CA 90806

The sipunculan fauna of southern California is poorly known. The first report was that of Chamberlain (1918). Fisher's (1950a,b, 1952) work provides the only major reference for this group. He erected subgenera for the genus *Golfingia* and described nearly all of the intertidal sipunculans, but only a few of the species from offshore. Recently however, several large scale surveys of the intertidal and benthic habitats of the entire southern California borderland have collected many species not reported from this area, including several apparently new taxa. This listing includes all previously reported sipunculans and also includes species seen in collections from southern California. It includes species from intertidal to bathyal habitats.

The classification used is based upon the families of Stephen and Edmonds (1972), and the subgenera of Fisher (1952), as revised by Stephen and Edmonds (1972), Cutler and Murina (1977), and Cutler (1979).

*Sipuncula* Stephen, 1965

*Sipunculoidea* Sedgwick, 1898

*Sipunculida* Pickford, 1947

*Sipunculidae* Baird, 1868

*Siphonosoma* Spengel, 1912

*Siphonosoma* (*Siphonosoma*) Fisher, 1950

*Siphonosoma* (*Siphonosoma*) *ingens* Fisher, 1950

*Sipunculus* Linnaeus, 1766

*Sipunculus nudus* Linnaeus, 1766

- Golfingiidae Stephen & Edmonds, 1972  
*Golfingia* Lankester, 1885  
*Golfingia* (*Apionsoma*) Sluiter, 1902, sensu Cutler  
*Golfingia* (*Apionsoma*) *capitata* (Gerould, 1913)  
     *Fisherana capitata* (Gerould), 1913  
*Golfingia* (*Apionsoma*) *misakiana* (Ikeda, 1904)  
     *Golfingia hespera* (Chamberlain, 1919)  
*Golfingia* (*Apionsoma*) *trichocephala* (Sluiter, 1902)  
*Golfingia* (*Golfingia*) Fisher, 1950  
*Golfingia* (*Golfingia*) *margaritacea* (Sars, 1851)  
*Golfingia* (*Nephasoma*) Pergament, 1940  
*Golfingia* (*Nephasoma*) *eremita* (Sars, 1851)  
*Golfingia* (*Nephasoma*) *laetmophilia* Fisher, 1952  
*Golfingia* (*Nephasoma*) *minuta* (Keferstein, 1863)  
*Golfingia* (*Nephasoma*) *nicolasi* Thompson (1980)  
*Golfingia* (*Nephasoma*) *pellucida* (Keferstein, 1865)  
*Golfingia* (*Thysanocardia*) Fisher, 1950  
T. nigra    { *Golfingia* (*Thysanocardia*) *catharinae* (Grube, 1868)  
               *Golfingia procera* (Möbius, 1875)  
*Golfingia* (*Thysanocardia*) *macginitieei* Fisher, 1952  
*Onchnesoma* Koren and Danielssen, 1875  
*Themiste* Gray, 1828  
*Themiste* *dyscrita* (Fisher, 1952)  
*Themiste* *perimeces* (Fisher, 1928)  
*Themiste* *pyroides* (Chamberlain, 1919)  
*Themiste* *zostericola* (Chamberlain, 1919)  
  
Phascolosomatidae Stephen & Edmonds, 1972  
*Phascolosoma* Leuckart, 1828  
*Phascolosoma* (*Phascolosoma*) Stephen & Edmonds, 1972  
*Phascolosoma* (*Phascolosoma*) *agassizii* Keferstein, 1867


A Preliminary, Artificial Key to  
the Sipunculans of Southern California

by

Bruce Thompson, SCCWRP


1. Musculature of the ~~body~~<sup>Trunk</sup> wall organized into longitudinal bands.....2
  - No longitudinal muscle bands.....4
2. Tentacles do not surround mouth but are dorsal to it; introvert hooks present.....Phascolosoma agassizii
  - Tentacles surround mouth; introvert hooks absent.....3
3. Introvert with sub-triangular scale-like papilla; skin on trunk organized into squares; tentacles on flaps of skin surrounding the mouth.....Sipunculus nudus
  - No scale-like papilla on introvert; tentacles are festooned in longitudinal rows.....Siphonosoma ingens  
forming chains
4. Contractile vessel with tubules or villi.....5
  - Contractile vessel without tubules or villi.....6
5. Villi on contractile vessel short; tentacles in longitudinal rows; no hooks.....Thysanocardia nigra
  - Long tubules on contractile vessel; tentacles dichotomously branched.....Themiste sp.
6. Single retractor muscle extends to posterior end of body; single nepridium.....Onchnesoma sp. A,
  - 2 or 4 retractor muscles; paired nepridia.....Golfingia sp....7
7. 2 retractor muscles.....8
  - 4 retractor muscles.....14
8. Hooks present an introvert.....9
  - Hooks absent from introvert.....12

9. Introvert slightly longer than trunk; tentacles reduced to small lobes.....Golfingia minuta  
- Introvert shorter than trunk; tentacles digitate.....10
10. Double circle of tentacles around mouth; introvert hooks slightly bent and awl like; yellow, wart-like papillae.....Golfingia pellucida  
- Single circle of tentacles around mouth, introvert hooks and papilla not as above.....11
11. Retractors insert in posterior 1/2 of trunk, rectal caecum present; posterior end of trunk dark brown .....Golfingia sp. 2  
- Retractors, insert in middle 1/3 of trunk; no rectal caecum.....Golfingia laetmophilia
12. Introvert shorter than trunk; posterior end of trunk extened to form a short "tail".....Golfingia sp. 1  
- Introvert longer than trunk; no posterior tail.....13
13. Introvert up to 9x longer than trunk; a 4-6 digitate tentacles.....G. nicolasi  
- Introvert less than 2x length of trunk; 20 digitate tentacles.....G. eremita
14. Tentacles surround mouth; spindle muscle not attached to posterior end on trunk; paired, single lobed nephridia  
...Sabogenuo...golfingia.....G. marginatus  
margaritacea  
- Tentacles, if present, do not surround mouth; but are dorsal to it; spindle muscle attached to posterior end of trunk, paired, bilobed nephridia.Sabogenuo...Aponeuma...15
15. Introvert shorter than trunk; ventral retractors originate from the posterior 1/2 of the trunk  
.....G. capitata  
- Introvert 5-10 times length of trunk.....16
16. Tentacles, hooks, papilla present.....G. misakiana  
- Tentacles, hooks, papilla, absent.....G. trichocephala




SIPUNCULIDAE

1. *Sipunculus*    2. *Xenosiphon*    3. *Siphonosoma*    4. *Phascolopsis*




GOLFINGIIDAE

5. *Golfingia*    6. *Onchnesoma*    7. *Themiste*    8. *Phascolion* (a. in shell, b. removed)



ASPIDOSIPHONIDAE

9. *idiosiphon*    10. *Paraspidosiphon*    11. *Cloeosiphon*  
12. *Lithacrosiphon*



PHASCOLOSMATIDAE

13. *Phascolosoma*

(From Edmonds, 1980; Stephen and Edmonds, 1972)

scale in mm

Characteristic Features of Families of the Phylum Sipuncula

Family	Integumentary canals or sacs in the body wall	Longitudinal musculature of the body wall	Arrangement of the tentacles	Anterior cap or shield	Genera
Sipunculidae	present (except in <i>Phascolopsis</i> )	grouped into bundles	numerous tentacles or a tentacular fold surround the mouth	absent	<i>Sipunculus</i> <i>Xenosiphon</i> <i>Siphonosoma</i> <i>Siphonomecus</i> <i>Phascolopsis</i>
Golfingiidae	absent	continuous	tentacles surround the mouth; may be in groups, be dendritic or reduced to a few lobes; may be absent in two genera	absent	<i>Golfingia</i> <i>Onchnesoma</i> <i>Nephasoma</i> <i>Phascolion</i> <i>Themiste</i>
Aspidosiphonidae	absent	either continuous or in bundles	surround or partly surround the mouth or in a ring dorsal to the mouth	present	<i>Aspidosiphon</i> <i>Paraspidosiphon</i> <i>Lithacrosiphon</i> <i>Cloeosiphon</i> <i>Centrosiphon</i>
Phascolosomatidae	absent (except in one species)	either continuous or in bundles	always in a horseshoe-shaped ring that is dorsal to the mouth; tentacles always enclose the nuchal organ,	absent	<i>Phascolosoma</i> <i>Fisherana</i>

## References

- Chamberlain, R.V. 1919. Notes on the Sipunculida of Laguna Beach. Pomona Coll. J. Ent. Zool. 12:30-31.
- Cutler, E.B. 1979. A reconsideration of the sipunculan taxa Fisherana Stephen, Mitosiphon Fisher, and Apionsoma Sluiter. Zool. Linn. J. Soc. London 65:367-384.
- Culter, E.B., and Murina, V.V. 1977. On the sipunculan genus Golfingia Lankester, 1885. Zool. J. Linn. Soc. London 60(2):173-187.
- Fisher, W.K. 1950. The sipunculid genus Phascolosoma. Ann. Mag. Nat. His. ser. 12, 3:547-552.
- Fisher, W.K. 1950. The new sub-genera and a new species of Siphonosoma (Spunculoidea). Ann. Mag. Nat. Hist. ser. 12, 3:805-808.
- Fisher, W.K. 1952. The sipunculid worms of California and Baja California. Proc. U.S. Nat. Mus. 102:371-450.
- Gibbs, P.E., E.B. Cutler, N.J. Cutler. 1983. A review of the Sipuncula genus Thypanocardia Fisher. Zool. Scripta, 12(4):295-304.
- Stephen, A.C. 1965. A revision of the phylum Sipuncula. Ann Mag. Nat. Hist. ser. 13, 7:457-462.
- Stephen, A.C. & S.I. Edmonds. 1972. The Phyla Sipuncula and Echiura. London: Trustees Brit. Mus. (Nat. Hist.), publ. No. 717, 528 pp.
- Thompson, B.E. 1980. Description of a new bathyal sipunculan from southern California, with ecological notes. Deepsea Research, 27(11):951-958.

2000-00000000

ангарын тохиолдсандаа энд нь залоги ЧИКИ .У.Я ,шилжүүлэг  
.ИЕ-00001 .Лодж .ЗСМ .Т .Нийгээний түншний

хувь нийтийн энд нь залжсанайландаа ЧИКИ .У.Я ,шилжүүлэг  
.ИЕ-00001 .Лодж .ЗСМ .Т .Нийгээний түншний түншний  
.ЧИКИ-САДЫГ нийтийн энд нь залоги .Лодж

хувь нийтийн энд нь залоги .У.Я ,шилжүүлэг .ЧИКИ  
.ИЕ-00001 .Лодж .ЗСМ .Т .Нийгээний түншний түншний  
.ЧИКИ-САДЫГ нийтийн энд нь залоги

хувь нийтийн энд нь залоги .ЧИКИ .У.Я ,шилжүүлэг  
.ИЕ-00001 .Лодж .ЗСМ .Т .Нийгээний түншний түншний

хувь нийтийн энд нь залоги .ЧИКИ .У.Я ,шилжүүлэг  
.ИЕ-00001 .Лодж .ЗСМ .Т .Нийгээний түншний түншний  
.ЧИКИ-САДЫГ нийтийн энд нь залоги

хувь нийтийн энд нь залоги .ЧИКИ .У.Я ,шилжүүлэг  
.ИЕ-00001 .Лодж .ЗСМ .Т .Нийгээний түншний түншний

хувь нийтийн энд нь залоги .ЧИКИ .У.Я ,шилжүүлэг  
.ИЕ-00001 .Лодж .ЗСМ .Т .Нийгээний түншний түншний

хувь нийтийн энд нь залоги .ЧИКИ .У.Я ,шилжүүлэг  
.ИЕ-00001 .Лодж .ЗСМ .Т .Нийгээний түншний түншний

хувь нийтийн энд нь залоги .ЧИКИ .У.Я ,шилжүүлэг  
.ИЕ-00001 .Лодж .ЗСМ .Т .Нийгээний түншний түншний

хувь нийтийн энд нь залоги .ЧИКИ .У.Я ,шилжүүлэг  
.ИЕ-00001 .Лодж .ЗСМ .Т .Нийгээний түншний түншний